

Guidance Document for Ordering and Usage of Trans- Isolate (T- I)



Toolkit

This is a guidance document that assists laboratorians with ordering, transportation and handling of T- I media.

TOOL CONTENTS:

- Guidance document for ordering and usage of Trans- Isolate (T- I)
 - T- I Media Questionnaire
-

ACKNOWLEDGEMENTS

This document is the product of collaborative efforts between several agencies, including the U.S. Centers for Disease Control and Prevention (CDC), the World Health Organization (WHO) Headquarters, WHO–Regional Office for Africa (WHO–AFRO), WHO–AFRO Inter–Country Support Teams (WHO–AFRO/IST), Agence de Médecine Préventive (AMP), Ministries of Health and the National Reference Laboratories from Burkina Faso, Mali, Niger, Chad and Togo, the Bill & Melinda Gates Foundation, Gavi, the Vaccine Alliance and the CDC Foundation. The authors thank all MenAfriNet Consortium members and members of the Laboratory Working Group for their input for the creation of the T- I guidance document.

For information about using the toolkit: MenAfriNet.org



Guidance Document for Ordering and Usage of Trans- Isolate (T- I)

I. What is T- I medium?

Trans- Isolate (T- I) is a biphasic medium used for the growth, holding, and transport of bacterial meningitis pathogens such as *Neisseria meningitidis*, *Haemophilus influenzae*, and *Streptococcus pneumoniae*. The solid phase of T- I is an agar slant that contains activated charcoal, soluble starch, and agar, while the liquid phase consists of a soy broth and supplement B (Figure 1). These supplements and nutrients will promote and support the growth of meningitis pathogens in both phases.



Figure 1.
Source: U.S. Centers for Disease Control and Prevention

II. Procuring T- I media from World Health Organization (WHO)

■ Estimate the quantity of T- I that needs to be ordered for the season:

- To estimate the needs for the upcoming season, the national reference laboratory (NRL) will need to average the total number of specimens received from across the country in the previous two years. A 10% contingency may be added to this number. Historical information on the number of specimens received per year at NRL can also be found in the WHO weekly meningitis bulletin. The actual amount to be ordered will be the difference between the needs for the season and the current stock inventory.
- Laboratories will need to get an accurate count of how much unexpired T- I media they have on inventory in order to determine how much T- I media they will need to order. It is important to exclude those media expiring before the end of the season.

■ Ordering T- I:

- The NRL will need to work with the Ministry of Health (MOH) to place the orders through the WHO Country Office.
- The country MOH will initiate the request with the WHO Country Office via email or letter.
- The WHO Country Office will forward the request to WHO Inter-Regional Support Team West Africa (ISTWA). The laboratory point of contact will need to be included in all correspondences.
- Once the order is processed, ISTWA will ship the media to the WHO Country Office, who will then transfer the order to the MOH.
- In order to improve ISTWA's production forecasts for the next season, a questionnaire will accompany the shipment (Appendix I). Laboratories are strongly encouraged to submit their feedback, as this will allow ISTWA to improve future forecasts.

III. T- I handling and transport

■ Quality control (QC)

- Non-inoculated T- I should be stored at 4°C immediately upon arrival in the laboratory.
- Remove T- I bottles from the refrigerator at least 30 minutes prior to inoculating it with the cerebrospinal fluid (CSF) specimen and allow it to warm to room temperature.
- NRL should perform quality control on all new shipments of T- I before use. For each new lot and shipment, perform and document the following:

- The following should be observed and documented:

All laboratories (NRL, peripheral)

- ♦ *Visual inspection:* Upon receipt, all media should be visually inspected for signs of contamination and physical defects at NRL and peripheral labs. Examples of contamination include turbid liquid, color change, or growth of bacteria or mold on the slant; physical defects include cracked glass, broken seal, leaking, reduced volume, or absent liquid phase. If any of the above occur, the T- I bottle should be discarded and ISTWA should be contacted for a replacement.

NRL and regional laboratories with culture capacity

- ♦ *Sterility testing:*
Incubate one T- I bottle vented and one non-vented (refer to T- I inoculation and venting procedure below) for 48 hours at 37°C.
 - » Using a sterile syringe, withdraw 50 µl of liquid phase of T- I and plate onto a chocolate agar plate.
 - » Incubate the plate at 37°C with 5% CO₂ for 48 hours. Any growth on the media is indicative of contamination.
- ♦ *Growth promotion:* Using sterile technique, inoculate one T- I bottle with 100 µl of an inoculum containing 10³ CFU/ml for each of the following pathogens: *N. meningitidis*, *H. influenzae* and *S. pneumoniae*. Incubate the plate at 37°C with 5% CO₂ for 48 hours and observe for growth of bacterial colonies.

■ T- I inoculation

- Prior to inoculation, disinfect the rubber stopper of the T- I bottle with 70% alcohol by first lifting the small metal cap on top of the T- I bottle. Do not completely remove the aluminum cover (Figure 2).
- Inoculate the T- I with 0.5 - 1.0 ml of cerebrospinal fluid (CSF) using 21G (0.88mm) sterile syringe through the rubber stopper of the lid.
- Invert the T- I bottle several time and incubate the inoculated T- I at 37°C.
- Immediately label the T- I bottle clearly with the date, name of the patient, and/or Epidemiological identification (EPID number) Country Code–Region –District–Year–Number (CCC- RRR- DDD- YY-NNNN). Include any other necessary information.



Figure 2.
Source: U.S. Centers for Disease Control and Prevention

■ Transport of T- I media

- If T- I media cannot be shipped within 24 hours, ventilate with a cotton plugged needle inserted through the rubber stopper without touching the media (Figure 3) and incubate at 37°C. Before transporting the T- I media, remove the venting needle, disinfect the rubber stopper, and transport in triple packaging following guidance for transport of infectious biological material. Transport T- I at ambient temperature.
- Dispose of all inoculated T- I in the same manner as infectious bacterial cultures.
- A case report form should be shipped along with the T- I to the reference laboratory as soon as possible.



Figure 3.
Source: U.S. Centers for Disease Control and Prevention

T-I Media Questionnaire

Appendix I.

Please answer all questions and email form back to the WHO-IST MDSC Point of Contact. Please do not leave any questions unanswered.

I. CURRENT STOCK

1. How many T- I media remain unused and in your possession? _____

When do these T- I media expire? _____ / _____ / _____ (dd/mm/yyyy)

II. NEW ORDER

2. How many T- I media have been requested? _____

What date did was the order submitted? _____ / _____ / _____ (dd/mm/yyyy)

When did the order arrive to the lab? _____ / _____ / _____ (dd/mm/yyyy)

How many T- I media were received by the laboratory? _____

What is the date of expiration of the T- I received? _____ / _____ / _____ (dd/mm/yyyy)

III. QUALITY CONTROL (QC)

3. Did you visually check all T- I media for contamination upon arrival? _____

4. What are the storage conditions of the T- I in your laboratory? _____

Stored at +2°C to +8°C

Stored at room temperature?

What temperature? _____

What precautions do you take before distributing the T- I to other laboratories? _____

5. Did you perform QC on the T- I?

Yes; How many? _____

No

6. Were there any issues with the QC?

No

Yes; Please describe: _____

IV. DISTRIBUTION TO LABORATORIES

7. How many T- I media do you keep at your lab? _____

8. How many T- I media were shared with or given to others? _____

Please provide the number and name of recipients. _____

9. Of those distributed to the laboratories, how many inoculated T- I did you receive back? _____

10. How are T- I distributed to from the National Reference Laboratory to peripheral labs?

In a cooler

At room temperature

By other means, please describe: _____

11. How are T- I transported from peripheral laboratories to National Reference Laboratory?

In a cooler

At room temperature

By other means, please describe: _____